



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY Speaking

Place your order with CEDARLANE® or your local distributor.  
Please contact CEDARLANE® for lot specific information.

## Mouse Anti-Human CD31 Monoclonal Antibody

CLSG36582  
CLSG36582-05  
Lot:

<b>Immunogen:</b>	BALB/C mice were immunized with membrane preparation of a spleen from a patient with hairy cell leukemia.
<b>Clone:</b>	JC/70A
<b>Isotype:</b>	IgG1, Kappa
<b>Format:</b>	1 ml (CLSG36582) or 0.5 ml (CLSG36582-05). This antibody is supplied as tissue culture supernatant containing sodium azide as preservative.
<b>Titer/Working Dilution:</b>	This antibody may be diluted to a titer of 1:10-1:25 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.
<b>Staining Protocol:</b>	We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. <u>Enzymatic digestion of paraffin embedded tissues is recommended prior to the immunostaining.</u>
<b>Specificity:</b>	This antibody reacts with a 100 kDa glycoprotein expressed by endothelial cells and 130 kDa glycoprotein present in platelets. It stains endothelial cells in normal as well as malignant tissues.
<b>Positive Control:</b>	Tonsil
<b>Cellular Localization:</b>	Cell membrane
<b>Storage:</b>	Store the antibody at 4-8°C. Do not use it beyond the expiration date stated on the label.
<b>References:</b>	i) Parums <i>et al.</i> J Clin Pathol 43: 752, 1990.

**Laboratory Reagent For Research Use Only**

JV 05/29/07

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



**toll free: 1-800-268-5058**  
**in North America**

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information  
website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)