



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY Speaking

Place your order with CEDARLANE® or your local distributor.
Please contact CEDARLANE® for lot specific information.

Purified Mouse Anti-Human CD44/HCAM Monoclonal Antibody

CLSG36595
CLSG36595-05
Lot: H892

Immunogen:	BALB/C mice were injected with stimulated human leukocytes.
Clone:	156-3C11
Isotype:	IgG2a
Format:	1 ml (CLSG36595) or 0.1 ml (CLSG36595-05). This antibody is supplied as purified immunoglobulin fraction containing sodium azide as preservative.
Titer/Working Dilution:	This antibody may be diluted to a titer of 1:25-1:75 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.
Staining Protocol:	We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. <u>Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.</u>
Specificity:	This antibody is specific to a cell surface glycoprotein of 80-95 kDa, known as CD44. CD44 is expressed on the surface of lymphocytes, monocytes and granulocytes. This antibody reacts with red cells pretreated with 2-aminoethylisothiuronium bromide (AET).
Positive Control:	Tonsil
Cellular Localization:	Cell membrane
Storage:	Store the antibody at 4-8°C. Do not use it beyond the expiration date stated on the label.
Reference:	i) Schlossman et al. Leukocyte Typing V, 1713, Oxford Univ. Press, 1995.

Laboratory Reagent For Research Use Only

JV 06/16/08

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com