



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

## Purified Mouse Anti-Human T Cell, CD45RO Monoclonal Antibody

CLSG36603

CLSG36603-05

Lot:

**Immunogen:**

BALB/C mice were immunized with IL-2 dependent T cell line, CA1.

**Clone:**

UCHL1

**Isotype:**

IgG2a, Kappa

**Format:**

1 ml (CLSG36603) or 0.5 ml (CLSG36603-05). This antibody is supplied as purified immunoglobulin containing sodium azide as preservative.

**Titer/Working Dilution:**

This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

**Staining Protocol:**

We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Suitable for formalin/paraffin tissues.

**Specificity:**

This antibody reacts with a 180 kDa glycoprotein of CD45 family, occurring on most thymocytes and activated T cells but only on the portion of the resting T cells. It reacts with most thymocytes, a subpopulation of resting cells within both the CD4 and CD8 subsets and mature activated T cells.

**Positive Control:**

Tonsil

**Cellular Localization:**

Cell membrane

**Storage:**

Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

**References:**

- i) Smith et al. Immunol 58: 63, 1986.
- ii) Terry et al. Immunol 64: 331, 1988.
- iii) Hall et al. J Clin Pathol 40: 151, 1987.

**Laboratory Reagent For Research Use Only**

JV 07/30/07

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



**toll free: 1-800-268-5058**

**in North America**

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)