



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Mouse Anti-Cytokeratin (45 46 56.4 kD) Monoclonal Antibody

CLSG36662
CLSG36662-05
Lot:

Immunogen: BALB/C mice were immunized with crude extract of splenic cells from a nude mouse engrafted with MCF-7 cells.

Clone: MNF116

Isotype: IgG1, kappa

Format: 1 ml (**CLSG36662**), or 0.5 ml (**CLSG36662-05**) This antibody is supplied as culture supernatant containing sodium azide as preservative.

Titer/Working Dilution: This antibody may be diluted to a titer of 1:25-1:50 using our PolyVue Detection System. The final dilution should be determined by the user based upon the staining conditions employed.

Staining Protocol: We suggest an incubation period of thirty minutes at room temperature. Optimal incubation conditions should be determined by the user based upon the fixation conditions and staining system employed. Enzymatic predigestion of paraffin embedded tissues is recommended prior to the immunostaining.

Specificity: This antibody recognizes cytokeratin polypeptides ranging from 40 through 58kD, corresponding to cytokeratins 5, 6, 8, 17, and 19. It shows a broad pattern of reactivity with human epithelial tissues from simple glandular epithelia to stratified squamous epithelia, like epidermis, mammary gland ducts, and tracheal epithelium.

Positive Control: Skin

Cellular Localization: Cytoplasmic

Storage: Store at 2-8°C. Do not use beyond the expiration date stated on the label.

References:
i) Gatter et al. J Clin Pathol 35: 1253, 1982.
ii) Gosh et al. Br J Haematol 61: 21, 1985.
iii) Norton et al. Histopathol 11: 487, 1987.

Laboratory Reagent For Research Use Only

BA 11/12/13

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: **Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: **Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com