



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)



Delivering You Today's Innovations for  
the Science of Tomorrow™

## Purified Mouse anti-Human MART-1/Melan-A Monoclonal Antibody

CLSG36774  
CLSG36774-05  
Lot:

**Immunogen:** BALB/C mice were injected with recombinant MART-1 protein.

**Clone:** A103

**Isotype:** IgG1

**Format:** 1 ml (CLSG36774), or 0.5 ml (CLSG36774-05). This antibody is supplied as purified immunoglobulin fraction containing sodium azide as a preservative.

**Titer/Working Dilution:** This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. The final dilution should be determined by the user based upon the staining conditions employed.

**Staining Protocol:** We suggest an incubation period of 30 minutes at room temperature. Optimal incubation conditions should be determined by the user based upon the fixation conditions and staining system employed. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

**Specificity:** This antibody is specific to a 20 kDa protein known as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. This antibody does not cross react with MAGE-1 or tyrosinase protein. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T-lymphocytes.

**Positive Control:** Melanoma

**Cellular localization:** Cytoplasmic

**Storage:** Store at 2-8°C. Do not use beyond the expiration date stated on the label.

**References:** i) Chen et al. Proc Natl Acad Sci USA 93: 5915, 1996.

**Laboratory Reagent For Research Use Only**

JV 09/01/10

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup>



[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)