



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Purified Mouse Anti-Human Cytokeratin 20 Monoclonal Antibody

CLSG36687

Lot:

Immunogen:	Cytokeratins isolated from human micro dissected villi of human duodenal mucosa.
Clone:	KS 20.8
Isotype:	IgG2a, kappa
Format:	1 ml of purified immunoglobulin containing sodium azide as preservative.
Titer/Working Dilution:	This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.
Staining Protocol:	We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. <u>Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.</u>
Specificity:	Cytokeratin 20 is less acidic than the other type I cytokeratins and is of particular interest because of its restricted range of expression. This antibody reacts with cytokeratin polypeptide of 46 kD in immunoblot analyses. Studies have shown that a marked difference exists in the expression of cytokeratin 20 among various carcinomas.
Positive Control:	Skin
Cellular Localization:	Cytoplasmic
Other applications:	Western Blotting (1:50 to 1:100)

Continued Overleaf...

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

Storage:

Store the antibody at 4-8°C. Do not use it beyond the expiration date stated on the label.

References:

- i) Angus et al. J Pathol 155: 71, 1988.
- ii) Gatter et al. J Clin Pathol 35: 1253, 1982.
- iii) Gosh et al. British J Hematol 153: 377, 1987.
- iv) Moll et al. Cell 31: 11, 1982.
- v) Takahashi et al. Path Res Pract 187: 145, 1991.

For Research Use Only. Not for Diagnostic or Therapeutic Work.

JV 05/03/06