



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)



Delivering You Today's Innovations for  
the Science of Tomorrow™

## Mouse anti-Human Hairy Cell Leukemia Monoclonal Antibody

CLSG36733  
CLSG36733-05  
Lot:

- Immunogen:** Deau cell line established from a large cell lymphoma.
- Clone:** DBA.44
- Isotype:** IgM, Kappa
- Format:** 1 ml (CLSG36733), or 0.5 ml (CLSG36733-05). This antibody is supplied as tissue culture supernatant containing sodium azide as a preservative.
- Titer/Working Dilution:** This antibody may be diluted to a titer of 1:15-1:30 in an ABC method. The final dilution should be determined by the user based upon the staining conditions employed.
- Staining Protocol:** We suggest an incubation period of 30 minutes at room temperature. Optimal incubation conditions should be determined by the user based upon the fixation conditions and staining system employed. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.
- Specificity:** This antibody reacts with a subset of B-lymphocytes localized in the follicular mantle zone. It reacts with 97% of hairy cell leukemia cases. This antibody shows strong positive staining of about 35% of cases of high grade B cell lymphomas.
- Positive Control:** Tonsil
- Cellular Localization:** Cell membrane, cytoplasmic
- Storage:** Store at 2-8°C. Do not use beyond the expiration date stated on the label.
- References:** i) Al Saati et al. Blood 74:2476, 1989.

**Laboratory Reagent For Research Use Only**

JV 07/16/10

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup>



[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

**In CANADA: Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

**In the USA: Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)