



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Purified Mouse Anti-Mast Cell Tryptase Monoclonal Antibody

CLSG36777
CLSG36777-05
Lot:

- Immunogen:** BALB/C mice were injected with a purified human lung Tryptase.
- Clone:** AA1
- Isotype:** IgG1
- Format:** 1 ml (CLSG36777) or 0.5 ml (CLSG36777-05). This antibody is supplied as purified immunoglobulin fraction containing sodium azide as preservative.
- Titer/Working Dilution:** This antibody may be diluted to a titre of 1:100-1:200 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.
- Staining Protocol:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the end user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.
- Specificity:** This antibody reacts with mast cells distributed in skin, synovium, lung and heart. This antibody does not bind with any other cell type. Mast cells contain a number of preformed chemicals mediators such as histamine, Chymase, carboxypeptidase and proteolytic Tryptase. Human mast cell tryptase is considered to be an important marker of mast cell activation as well as an important mediator of inflammation. This antibody cross-reacts with monkey, dog and cat.

Continued Overleaf...

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

Positive Control:

Tonsil

Cellular Localization:

Cytoplasmic

Other Applications:

Western Blotting

Storage:

Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

References:

i) Walls et al. Clin Exp Allergy 20: 581, 1990.

ii) Walls et al. J Pathol 162: 119, 1990.

Laboratory Reagent For Research Use Only

JV 07/03/07