



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Delivering You Today's Innovations for
the Science of Tomorrow™

Purified Mouse Anti-MyoD1 Monoclonal Antibody

CLSG36792
CLSG36792-05

Lot:

Immunogen: BALB/C mice were injected with a recombinant mouse MyoD1 protein.

Clone: 5.2F

Isotype: IgG2a, kappa

Format: 1 ml (**CLSG36792**) or 0.5 ml (**CLSG36792-05**). This antibody is supplied as purified immunoglobulin fraction containing sodium azide as preservative.

Titer/Working Dilution: This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

Staining Protocol: We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 1mM EDTA, pH 8 or 10 mM citrate buffer, pH 6.0 prior to immunostaining.

Specificity: This antibody is specific to a 45 kDa protein, which is identified as MyoD1. This antibody is specific to an epitope of amino acid 3-56 in mouse MyoD1. It does not cross-react with myogenin, Myf5 or Myf6. This antibody stains the nuclei of myoblasts in developing muscle tissue. MyoD1 is not detected in normal adult tissue but is expressed strongly in the tumor cell nuclei of rhabdomyosarcomas. This antibody cross-reacts with human, mouse, rat and chicken.

Positive Control: Rhabdomyosarcoma

Cellular Localization: Nuclear

Storage: Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

References:

1. Dias et al. Cancer Res 52: 144, 1994.
2. Rosai et al. Am J Surg. Pathol. 15: 974, 1991.
3. Tallini et al Am J Pathol 144: 693, 1994.
4. Thulasi et al. Cell Growth Diff 7(4):531, 1996.

Laboratory Reagent For Research Use Only

JV 09/17/09

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com