



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Mouse Anti-Human Proliferating Cell Nuclear Antigen (PCNA) Monoclonal Antibody

CLSG36826

CLSG36826-05

Lot:

Immunogen:

Rat PCNA made in the protein A expression vector pR1T2T.

Clone:

PC10

Isotype:

IgG2a, Kappa

Format:

1 ml (CLSG36826) or 0.5 ml (CLSG36826-05). This antibody is supplied as diluted ascites containing sodium azide as preservative.

Titer/Working Dilution:

This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

Staining Protocol:

We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

Specificity:

PCNA is essential for cellular DNA synthesis. This antibody reacts with proliferating cells in a wide range of normal tissues. The significance of PCNA immunoreactivity in tumors is not fully established yet. However, in non-Hodgkin's lymphoma a linear relation between Ki-67 and PCNA staining has been demonstrated.

Positive Control:

Tonsil

Cellular Localization:

Nuclear

Storage:

Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

References:

- i) Waseem et al. J Cell Sci 96: 121, 1990.
- ii) Takasaki et al. J Exp Med 159: 981, 1984.
- iii) Hall et al. J Pathol 162: 285, 1990.
- iv) Coltrera et al. Applied Immunohistochemistry 1: 193, 1993.

Laboratory Reagent For Research Use Only

JV 05/06/08

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com