



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Purified Mouse Anti-Human Perforin Monoclonal Antibody

CLSG36827
CLSG36827-05

Immunogen:	Recombinant protein corresponding to an external region of the C-terminus of the perforin molecule.
Clone:	5B10
Isotype:	IgG1
Format:	1 ml (CLSG36827) or 0.5 ml (CLSG36827-05). This antibody is supplied as purified immunoglobulin fraction containing <0.1% NaN ₃ as preservative.
Titer/Working Dilution:	This antibody may be diluted to a titer of 1:10-1:20 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.
Staining Protocol:	We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the end user. <u>Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.</u>
Specificity:	This antibody is specific to a 70 kDa protein, which is a potent cytolytic pore-forming protein. Perforin is a specific marker of functionally active cytotoxic T-lymphocytes and natural killer cells. Perforin mediates cytolysis of target cells by membrane damage and apoptosis.
Positive Control:	Spleen
Cellular Localization:	Cytoplasmic
Storage:	Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.
References:	i) Hameed et al. Am J Pathol 140 (5): 1025, 1992. ii) Lichtenheld MG et al. Nature 335: 448, 1988.

Laboratory Reagent For Research Use Only.

JK 03/22/17

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com