



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY Speaking

Place your order with CEDARLANE® or your local distributor.
Please contact CEDARLANE® for lot specific information.

Mouse Anti-Human TAG-72/CA 72-4 Monoclonal Antibody

CLSG36872
CLSG36872-05
Lot: H983

| | |
|--------------------------------|---|
| Immunogen: | BALB/C mice were injected with membrane enriched fraction of a human breast carcinoma liver metastasis. |
| Clone: | B72.3 |
| Isotype: | IgG1, kappa |
| Format: | 1 ml (CLSG36872) or 0.5 ml (CLSG36872-05). This antibody is supplied as tissue culture supernatant containing sodium azide as preservative. |
| Titer/Working Dilution: | This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. |
| Staining Protocol: | We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. <u>Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.</u> |
| Specificity: | This antibody is specific to an oncofetal antigen of > 1,000 kDa known as tumor associated glycoprotein (TAG-72) or CA 72-4. This antibody defines the mucin carried sialylated-Tn antigen. This antibody stains majority of human adenocarcinomas including colorectal, pancreatic, gastric, ovarian, endometrial, mammary and non-small cell lung cancer. |
| Positive Control: | Breast carcinoma |
| Cellular Localization: | Cytoplasmic, cell surface |
| Storage: | Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label. |

Laboratory Reagent For Research Use Only

JV 09/22/08

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com