



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Mouse Anti-Human Wilms' Tumor 1 Protein (WT1)

CLSG368781

CLSG368781-05

Lot:

Clone:

6F-H2

Immuogen:

BALB/C mice were injected with truncated human WT1 protein corresponding to N-terminal amino acids 1-181.

Isotype:

IgG₁

Format:

1 ml (CLSG368781) or 0.5 ml (CLSG368781-05). This antibody is supplied as tissue culture supernatant containing BSA and 0.05% sodium azide (NaN₃) as preservative.

Titer/Working

Dilution:

This antibody may be diluted to a titer of 1:15-1:25 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

Staining Protocol:

We suggest an incubation period of 30 to 60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM Citrate buffer, pH 6.0 prior to immunostaining.

Specificity:

Wilms' Tumor (WT1), is a gene involved in the induction of Wilms' tumor, a pediatric renal malignancy. Wilms' tumor is associated with mutations of WT1, a zinc-finger transcription factor that is essential for the development of the metanephric kidney and the urogenital system. The WT1 gene is normally expressed in fetal kidney and mesothelium, and its expression has been suggested as a marker for Wilms' tumor and mesothelioma.

Positive Control:

Wilms' tumor

Cellular Localization:

Nuclear and/or Cytoplasmic

Storage:

Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

References:

- i) Haber et al. New Biol 4: 97, 1992.
- ii) Kumar-Singh et al. J Pathol 181: 67, 1997.
- iii) Menssen et al. Leukemia 9: 1060, 1995.

Laboratory Reagent For Research Use Only

JV 10/04/07

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058

in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com