



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Purified Rabbit anti-CD3 Zeta Polyclonal Antibody

CLSG36906

CLSG36906-05

Lot:

Immunogen:

A synthetic peptide corresponding to a.a. 36 –54 a.a. of the human CD3 zeta protein.

Host:

Rabbit

Format:

1 ml (**CLSG36906**) or 0.5 ml (**CLSG36906-05**). Purified immunoglobulin fraction of rabbit antiserum against CD3 zeta containing sodium azide as preservative.

Titer/Working Dilution:

This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

Staining Protocol:

We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

Specificity:

This antibody reacts with a 19 kDa protein. CD3 consists of five invariant polypeptide chains that associate to form three dimers: a heterodimer of gamma and epsilon chains, a heterodimer of delta and epsilon chains and a homodimer of two zeta chains or a heterodimer of zeta and eta chains. Zeta chain consists of three consecutive copies of the same motif. This antibody cross reacts with mouse and rat.

Positive Control:

Tonsil

Cellular Localization:

Cell membrane

Storage:

Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

Laboratory Reagent For Research Use Only

JV 11/20/08

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058

in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com