



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY Speaking

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

## Purified Rabbit Anti-Glial Fibrillary Acidic Protein (GFAP)

**CLSG36937**

**Lot:** D943

**Immunogen:**

Glial fibrillary protein isolated from cow spinal cord.

**Host:**

Rabbit

**Size:**

1 ml

**Format:**

Purified immunoglobulin fraction of rabbit antiserum against GFAP containing sodium azide as preservative.

**Titer/Working Dilution:**

This antibody may be diluted to a titer of 1:50-1:250 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

**Staining Protocol:**

We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions should be determined by the user.

Suitable for formalin/paraffin tissues.

**Specificity:**

This antibody reacts with human GFAP as well as GFAP from cat, chicken, cow, guinea pig, hamster, monkey, rat, mouse, sheep and kangaroo. In the central nervous system, this antibody stains astrocytes and some groups of ependymal cells. However, in the peripheral nervous system, it stains Schwann cells, satellite cells and enteric glial cells.

**Positive Control:**

Brain

**Cellular Localization:**

Cytoplasmic

**Storage:**

Store the antibody at 4-8°C. Do not use it beyond the expiration date stated on the label.

**Reference:**

1. Viale *et al.* Arch. A. Pathol. Anat. **418**:339, 1991.

**FOR RESEARCH USE ONLY**

CH 08/23/05

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



**toll free: 1-800-268-5058**  
**in North America**

**phone: (905) 878-8891 • fax: (905) 878-7800**

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information  
**website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)**