



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.  
Please contact CEDARLANE® for lot specific information.

## Purified Rabbit Anti-Human IgE Polyclonal Antibody

CLSG36950  
CLSG36950-05  
Lot:

- Immunogen:** IgE isolated from the serum of a patient with IgE-myeloma.
- Host:** Rabbit
- Format:** 1 ml (CLSG36950) or 0.5 ml (CLSG36950-05), purified immunoglobulin fraction of rabbit antiserum against human IgE containing sodium azide as preservative.
- Titer/Working Dilution:** This antibody may be diluted to a titer of 1:50-1:500 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.
- Staining Protocol:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions should be determined by the user. Proteolytic treatment is required prior to immunostaining of formalin fixed paraffin embedded tissue sections.
- Specificity:** This antibody reacts with the epsilon chain of human IgE. Traces of contaminating antibodies have been removed by solid phase absorption. Only the IgE arch appeared in crossed immunoelectrophoresis.
- Positive Control:** Tonsil
- Cellular Localization:** Cytoplasmic
- Storage:** Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

**Laboratory Reagent For Research Use Only.**

JV 08/24/07

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



*toll free: 1-800-268-5058*  
*in North America*

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information  
website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)