



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>   
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

## Rabbit Polyclonal Antibody to Human Pancreatic Polypeptide

CLSG36973, 1 ml  
CLSG36973-05, 0.5 ml  
Lot: S113

<b>Intended Use:</b>	This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy. Interpretation of staining results should be accompanied by histological studies with proper controls.
<b>Immunogen:</b>	Synthetic human pancreatic polypeptide conjugated to keyhole limpet hemocyanin.
<b>Host:</b>	Rabbit
<b>Format:</b>	Purified immunoglobulin fraction against human pancreatic polypeptide containing sodium azide as a preservative.
<b>Titer/Working Dilution:</b>	This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. The final dilution should be determined by the user based upon the staining conditions employed.
<b>Staining Protocol:</b>	We suggest an incubation period of 30 minutes at room Temperature. Optimal incubation conditions should be determined by the user based upon the fixation conditions and staining system employed. Suitable for formalin fixed, paraffin embedded tissue sections.
<b>Specificity:</b>	This antibody reacts with pancreatic polypeptide. It stains the periphery of islets, exocrine pancreatic parenchyma, and the epithelium of small and medium sized ducts and acinar cells.
<b>Positive Control:</b>	Pancreas
<b>Cellular Localization:</b>	Cytoplasmic
<b>Storage:</b>	Store at 2-8°C. Do not use beyond the expiration date stated on the label.

**Laboratory Reagent For Research Use Only**

EJ 11/11

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup> 

[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)