



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Purified Rabbit Anti-PARP (Poly ADP-Ribose Polymerase) Polyclonal Antibody

CLSG36975

CLSG36975-05

Lot:

Immunogen:

A synthetic peptide from the N-terminal region of human PARP.

Host:

Rabbit

Format:

1 ml (CLSG36975) or 0.5 ml (CLSG36975-05). Purified immunoglobulin fraction of rabbit antiserum against human PARP containing sodium azide as preservative.

Titer/Working Dilution:

This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

Staining Protocol:

We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions should be determined by the user. High temperature treatment of formalin-fixed tissue sections with 10 mM citrate buffer, pH 6.0 must be performed prior to the immunostaining.

Specificity:

This antibody reacts with a 116 kDa protein, known as PARP. Proteolysis of PARP to its stable 85 kDa fragment is an early marker of apoptosis and is mediated by Caspase (CPP32) protein. PARP uses nicotinamide adenine dinucleotide (oxidized form) as a substrate to catalyze the transfer of ADP-ribose to a variety of nuclear acceptors. This antibody cross reacts with human, mouse and rat.

Positive Control:

Tonsil

Cellular Localization:

Nuclear

Other Applications:

Western Blotting

Storage:

Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

References:

- i) Lamarre et al. Biochimica et Biophysica Acta 950: 147, 1988.
- ii) Tewari et al. Cell 81: 801, 1995.

Laboratory Reagent For Research Use Only

JV 04/07/08

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058

in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com