



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Purified Mouse Anti-Human IgE Monoclonal Antibody

CLX166AP

Lot:

Clone: 4G7

Isotype: Mouse IgG

Specificity: The antibody 4G7 reacts with human IgE; it recognizes an epitope different from the ones recognized by BE5 and 4H10 antibodies to IgE.

Immunogen: Purified human IgE.

Species Reactivity: Human

Application: ELISA

Purity: > 95% (by SDS-PAGE).

Purification: Purified from ascites by precipitation methods and ion exchange chromatography.

Concentration: 0.1 mg at 1 mg/ml.

Storage Buffer: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4.

Storage / Stability: Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.

Background: Immunoglobulin E (IgE) is a 180 kDa soluble protein serving as an antigen-specific unit of mast cell effector mechanisms. IgE has the lowest serum concentration of all immunoglobulins (approximately 0.5 mg/l) in healthy individuals, but upon allergen challenge its concentration in blood increases dramatically. Although biological survival of free IgE is very short ($T_{1/2} = 2$ days), it is stabilized after binding to its high affinity receptor. Unlike IgM- IgG- and IgA-committed B cells, IgE-switched B cells do not undergo clonal expansion.

Continued Overleaf.....

Visit our website for your local distributor.

CEDARLANE®

www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.



In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

- References:**
- *Franklin EC: Structure and function of immunoglobulins. *Acta Endocrinol Suppl (Copenh)*. 1975;194:77-95.
 - *Fuller JM, Keyser JW: Serum immunoglobulins after surgical operation. *Clin Chem*. 1975 May;21(6):667-71.
 - *Balogh Z, Merétey K, Falus A, Bozsóky S: Serological abnormalities in juvenile chronic arthritis: a review of 46 cases. *Ann Rheum Dis*. 1980 Apr;39(2):129-34.
 - *Brinkmann V, Heusser CH: T cell-dependent differentiation of human B cells into IgM, IgG, IgA, or IgE plasma cells: high rate of antibody production by IgE plasma cells, but limited clonal expansion of IgE precursors. *Cell Immunol*. 1993 Dec;152(2):323-32.
 - *Gould HJ, Beavil RL, Vercelli D: IgE isotype determination: epsilon-germline gene transcription, DNA recombination and B-cell differentiation. *Br Med Bull*. 2000;56(4):908-24.
 - *Mueller GA, Smith AM, Chapman MD, Rule GS, Benjamin DC: Hydrogen exchange nuclear magnetic resonance spectroscopy mapping of antibody epitopes on the house dust mite allergen Der p 2. *J Biol Chem*. 2001 Mar 23;276(12):9359-65.
 - *Smith AM, Benjamin DC, Hozic N, Derewenda U, Smith WA, Thomas WR, Gafvelin G, van Hage-Hamsten M, Chapman MD: The molecular basis of antigenic cross-reactivity between the group 2 mite allergens. *J Allergy Clin Immunol*. 2001 Jun;107(6):977-84.

Laboratory Reagent For Research Use Only