



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Purified Mouse anti- Human CD11b Monoclonal Antibody

CLX17AP

Lot:

Size: 0.1 mg

Clone: MEM-174

Isotype: Mouse IgG2a

Specificity: The antibody MEM-174 recognizes CD11b antigen (Mac-1), a 165 kDa transmembrane protein type 1 mainly expressed on monocytes, granulocytes and NK-cells. The CD11b mediates neutrophil and monocyte interactions with stimulated endothelium.

HLDA VI; WS Code BP 310

HLDA VI; WS Code M 18

Immunogen: Human granulocytes.

Species Reactivity: Human

Continued Overleaf...

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

Application:	<p>Flow Cytometry <i>Positive control:</i> Peripheral Blood Lymphocytes</p> <p>Immunoprecipitation <i>Positive control:</i> Granulocytes CD11b/CD18 transfectants <i>Application note:</i> Immunoprecipitation with the antibody MEM-174 was successfully performed using immuno-affinity sorbents (minicolumns). The final immunosorbent should contain 1-5 mg of antibody/1 ml of sorbent gel (e.g. agarose beads) It is recommended to stimulate positive control material by LPS 1-2 days before to achieve better expression of target antigen).</p> <p>Functional Application The antibody blocks binding of LFA-1 complex to ICAM-1.</p>
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified from ascites by protein-A affinity chromatography.
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not use after expiration date stamped on vial label. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.
References:	Leukocyte Typing VI. Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).

This product is made available in collaboration with Exbio Praha, a.s.

For Research Use Only.