



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Purified Mouse anti-EGFR (Phospho-Tyr992) Monoclonal Antibody

CLX297AP

Lot:

Size: 0.1 mg

Concentration: 1 mg/ml

Clone: EM-12

Isotype: Mouse IgG1

Specificity: The antibody EM-12 reacts with human EGFR (ErbB1 / HER1) phosphorylated on tyrosine 992.

Immunogen: Synthetic phospho-peptide covering sequence around tyrosine 992 of human EGFR.

Species Reactivity: Human

Application:
Flow Cytometry
Immunoprecipitation
Western Blotting
Recommended dilution: 1 µg/ml
Positive control: EGF-stimulated A431
Negative control: Non-stimulated A431

Purity: > 95% (by SDS-PAGE)

Purification: Purified from hybridoma culture supernatant by protein-A affinity chromatography.

Storage Buffer: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4

Storage / Stability: Store at 2-8°C. Do not use after expiration date stamped on vial label. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.

Background: The oncoprotein EGFR (epidermal growth factor receptor), also known as HER1 / ErbB1, is a member of ErbB family of cell surface receptor tyrosine kinases. This 170 kDa transmembrane glycoprotein is often associated with cancerogenesis, although its activation state is controlled at various levels including trafficking and degradation steps. Binding of receptor-specific ligands to the EGFR ectodomain results in formation of homodimeric or heterodimeric kinase-active complexes into which HER2 / ErbB2 is recruited as a preferred partner. Subsequent signaling cascades such as RAS/MAPK and PI3K/AKT pathways lead to cell proliferation and survival responses.

Continued...

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

References:

- *Moulder SL, Yakes FM, Muthuswamy SK, Bianco R, Simpson JF, Arteaga CL: Epidermal growth factor receptor (HER1) tyrosine kinase inhibitor ZD1839 (Iressa) inhibits HER2/neu (erbB2)-overexpressing breast cancer cells in vitro and in vivo. *Cancer Res.* 2001 Dec 15;61(24):8887-95.
- *Batzer AG, Blaikie P, Nelson K, Schlessinger J, Margolis B: The phosphotyrosine interaction domain of Shc binds an LXNPXY motif on the epidermal growth factor receptor. *Mol Cell Biol.* 1995 Aug;15(8):4403-9.
- *Daniele T, Di Tullio G, Santoro M, Turacchio G, De Matteis MA: ARAP1 regulates EGF receptor trafficking and signalling. *Traffic.* 2008 Dec;9(12):2221-35.
- *Rappl A, Piontek G, Schlegel J: EGFR-dependent migration of glial cells is mediated by reorganisation of N-cadherin. *J Cell Sci.* 2008 Dec 15;121(Pt 24):4089-97.
- *Gandhi J, Zhang J, Xie Y, Soh J, Shigematsu H, Zhang W, Yamamoto H, Peyton M, Girard L, Lockwood WW, Lam WL, Varella-Garcia M, Minna JD, Gazdar AF: Alterations in genes of the EGFR signaling pathway and their relationship to EGFR tyrosine kinase inhibitor sensitivity in lung cancer cell lines. *PLoS ONE.* 2009;4(2):e4576.
- *Yan F, Cao H, Chaturvedi R, Krishna U, Hobbs SS, Dempsey PJ, Peek RM Jr, Cover TL, Washington MK, Wilson KT, Polk DB: Epidermal growth factor receptor activation protects gastric epithelial cells from Helicobacter pylori-induced apoptosis. *Gastroenterology.* 2009 Apr;136(4):1297-1307, e1-3.

Laboratory Reagent For Research Use Only

BA 06/04/13