



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Mouse Anti-CD203c Monoclonal Antibody

CLX310AP
CLX310F
CLX310PE
CLX310APC

Clone: NP4D6

Isotype: Mouse IgG1

Specificity:

The mouse monoclonal antibody NP4D6 reacts with CD203c, a transmembrane ectoenzyme expressed on basophils and mast cells, and overexpressed upon their activation.
HLDA VIII

Immunogen: HEK-293 cells transfected with human CD203c.

Species Reactivity: Human

Application: Flow Cytometry; Immunocytochemistry.

Conjugate Preparation:

The purified antibody is conjugated with Fluorescein isothiocyanate (FITC), R-Phycoerythrin (PE) or cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.

Presentation:

Purified: 0.1 mg (1 mg/mL) purified IgG buffered in PBS with 15 mM sodium azide, approx. pH 7.4. (Purified from hybridoma culture supernatant by protein-A affinity chromatography).

FITC: 0.4 mL of FITC conjugated IgG buffered in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. Sufficient for 100 tests.

PE: 2 mL of PE conjugated IgG buffered in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. Sufficient for 100 tests.

APC: 1 mL of APC conjugated IgG buffered in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. Sufficient for 100 tests.

Storage / Stability:

Store in the dark at 2-8°C. Do not freeze all formats. Avoid prolonged exposure to light of conjugates.

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2015 and ISO 13485:2016
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

Usage:

Recommended dilutions for Flow Cytometry analysis of human blood cells using:

Purified: 1-5 µg/ml

FITC: 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.

PE: 20 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.

APC: 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.

Background:

CD203c, also known as ENPP-3, is integral membrane ectoenzyme (ectonucleotide pyrophosphatase/phosphodiesterase 3), that hydrolyses nucleotide triphosphates and thus modulates purinergic signaling. CD203c is expressed mainly on activated basophils and mast cells. CD203c is upregulated in response to IgE-receptor cross-linking and is overexpressed on neoplastic mast cells in patients with systemic mastocytosis. Measurement of its induced enhancement on the plasma membrane is useful for diagnostics of allergies.

References:

- *Hauswirth AW, Escribano L, Prados A, Nuñez R, Mirkina I, Kneidinger M, Florian S, Sonneck K, Vales A, Scherthaner GH, Sanchez-Muñoz L, Sperr WR, Bühring HJ, Orfao A, Valent P: CD203c is overexpressed on neoplastic mast cells in systemic mastocytosis and is upregulated upon IgE receptor cross-linking. *Int J Immunopathol Pharmacol.* 2008 Oct-Dec;21(4):797-806.
- *Tokuda R, Nagao M, Hiraguchi Y, Hosoki K, Matsuda T, Kouno K, Morita E, Fujisawa T: Antigen-induced expression of CD203c on basophils predicts IgE-mediated wheat allergy. *Allergol Int.* 2009 Jun;58(2):193-9.
- *Wanich N, Nowak-Wegrzyn A, Sampson HA, Shreffler WG: Allergen-specific basophil suppression associated with clinical tolerance in patients with milk allergy. *J Allergy Clin Immunol.* 2009 Apr;123(4):789-94.e20.
- *Chirumbolo S, Brizzi M, Ortolani R, Vella A, Bellavite P: Inhibition of CD203c membrane up-regulation in human basophils by high dilutions of histamine: a controlled replication study. *Inflamm Res.* 2009 Nov;58(11):755-64.
- *Bühring HJ, Simmons PJ, Pudney M, Müller R, Jarrossay D, van Agthoven A, Willheim M, Brugger W, Valent P, Kanz L: The monoclonal antibody 97A6 defines a novel surface antigen expressed on human basophils and their multipotent and unipotent progenitors. *Blood.* 1999 Oct 1;94(7):2343-56.
- *Bühring HJ, Seiffert M, Giesert C, Marxer A, Kanz L, Valent P, Sano K: The basophil activation marker defined by antibody 97A6 is identical to the ectonucleotide pyrophosphatase/phosphodiesterase 3. 1. *Blood.* 2001 May 15;97(10):3303-5.
- *Platz IJ, Binder M, Marxer A, Lischka G, Valent P, Bühring HJ: Hymenoptera-venom-induced upregulation of the basophil activation marker ecto-nucleotide pyrophosphatase/phosphodiesterase 3 in sensitized individuals. *Int Arch Allergy Immunol.* 2001 Dec;126(4):335-42.
- *Heneberg P, Riegerová K, Kučera P: Pimecrolimus Is a Potent Inhibitor of Allergic Reactions to Hymenopteran Venom Extracts and Birch Pollen Allergen In Vitro. *PLoS One.* 2015 Nov 12;10(11):e0142953.

Laboratory Reagent For Research Use Only