



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Purified Mouse Anti-CD195 / CCR5 Monoclonal Antibody

CLX486AP

Size: 0.1 mg

Clone: T21/8

Isotype: Mouse IgG1

Specificity: The mouse monoclonal antibody T21/8 recognizes the N-terminus of CD195, an approximately 45 kDa G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.

Immunogen: CCR5 peptide (Met1-Lys22) KLH conjugate

Species Reactivity: Human

Application: Flow Cytometry
Immunoprecipitation
Western Blotting
Application note: do not boil cell lysates prior to SDS-PAGE
Immunocytochemistry
ELISA

Purity: > 95% (by SDS-PAGE)

Purification: Purified from cell culture supernatant by protein-A affinity chromatography.

Concentration: 1 mg/ml

Storage Buffer: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4.

Storage / Stability: Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.

Background: **CD195 / CCR5** (also known as CKR-5) is a receptor for inflammatory CC-chemokines (characterized by a pair of adjacent cysteine residues), such as MIP-1 alpha, MIP-1 beta, or RANTES. It is a G protein-associated seven-pass transmembrane protein expressed on resting T cells with memory/effector phenotype, monocytes, macrophages and immature dendritic cells. This chemokine receptor regulates the activation and directed migration of leukocytes. Importantly, along with CD4, CD195 / CCR5 functions as a major receptor for HIV. Their ligand is the viral glycoprotein gp120.

Continued Overleaf.....

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

References:

- *Pollok-Kopp B, Schwarze K, Baradari VK, Oppermann M: Analysis of ligand-stimulated CC chemokine receptor 5 (CCR5) phosphorylation in intact cells using phosphosite-specific antibodies. *J Biol Chem.* 2003 Jan 24;278(4):2190-8.
- *Hüttenrauch F, Pollok-Kopp B, Oppermann M: G protein-coupled receptor kinases promote phosphorylation and beta-arrestin-mediated internalization of CCR5 homo- and hetero-oligomers. *J Biol Chem.* 2005 Nov 11;280(45):37503-15.
- *Monde K, Maeda Y, Tanaka Y, Harada S, Yusa K: Gp120 V3-dependent impairment of R5 HIV-1 infectivity due to virion-incorporated CCR5. *J Biol Chem.* 2007 Dec 21;282(51):36923-32.
- *Mascalchi P, Lamort AS, Salomé L, Dumas F: Single Particle Tracking reveals two distinct environments for CD4 receptors at the surface of living T lymphocytes. *Biochem Biophys Res Commun.* 2012 Jan 6;417(1):409-13.

Laboratory Reagent For Research Use Only

JK 07/19/18