



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Delivering You Today's Innovations for
the Science of Tomorrow™

Human Tumor Necrosis Factor-alpha (AF)

CLY100-111AF-10UG
CLY100-111AF-50UG
CLY100-111AF-100UG
CLY100-111AF-500UG
CLY100-111AF-1MG

Description: Tumor Necrosis Factor alpha (TNF α) is an inflammatory cytokine secreted by macrophages, monocytes, neutrophils, T cells, NK-cells following their stimulation by bacterial LPS. TNF α activates signals through two receptors, TNFR1, which is expressed on most cell types, and TNFR2, which is expressed mainly on immune cells. TNF α can have many functions including, to stimulate phagocytosis in macrophages, to chemoattract neutrophils, to increase insulin resistance and to induce fever. Recombinant human TNF- α is a non-glycosylated protein, containing 158 amino acids, with a molecular weight of 17.5 kDa.

Source: *E.coli*. This product is produced with no animal or human origin raw products. All processing and handling employs animal free equipment and animal free protocols.

Presentation: 10 μ g (CLY100-111AF-10UG), 50 μ g (CLY100-111AF-50UG), 100 μ g (CLY100-111AF-100UG), 500 μ g (CLY100-111AF-500UG) or 1 mg (CLY100-111AF-1MG), sterile filtered white lyophilized (freeze-dried) powder. Recombinant human TNF- α is lyophilized from a concentrated solution (1mg/mL) with 10 mM Na₂PO₄, pH 8.0.

Reconstitution: Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/mL, which can be further diluted into other aqueous solutions.

Stability: Lyophilized product is very stable at -20°C. Reconstituted material should be aliquoted and frozen at -20°C. It is recommended that a carrier protein (0.1% HSA or BSA) is added for long term storage.

Protein Content and Purity (typically > 96%) determined by:
Reducing and non-reducing SDS-PAGE.

Endotoxin Level: Endotoxin level as measured by LAL is 0.05 EUs/ μ g.

Biological Activity: Recombinant Tumor Necrosis Factor-alpha (AF) is fully biologically active when compared to standard. The ED₅₀ as determined by the cytolysis of mouse L929 cells in the presence of Actinomycin D is typically less than 0.05 ng/ml.

AA Sequence: MVRSSSRTPS DKPVAHVVAN PQAEGQLQWL NRRANALLAN GVLRDNQLV
VPSEGLYLIY SQVLFKGGQC PSTHVLLTHT ISRIAVSYQT KVNLLSAIKS PCQRETPEGA EAKPWYEPIY
LGGVVFQLEKG DRLSAEINRP DYLDFAESGQ VYFGIIL

Laboratory Reagent For Research Use Only

MW 07/19/19

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com