

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Conveniently Delivering You Today's Innovations for the Science of Tomorrow™

Human Pleiotrophin

CLY100-55-5UG CLY100-55-20UG CLY100-55-100UG CLY100-55-500UG **CLY100-55-1MG**

Description: Pleiotrophin (PTN) is a heparin-binding growth factor that has mitogenic effects on fibroblast, epithelial, and endothelial cells. PTN is made by many tissues, but is predominantly secreted by nervous tissue during development. PTN induces neurite outgrowth and is involved in tumor growth and metastasis. PTN binds with low affinity to the cell surface receptor nucleolin to inhibit HIV-1 infection. PNT also binds the receptor protein tyrosine phosphatase type Z (PTPRZ), syndecan-3, and anaplastic lymphoma kinase (ALK) receptors.

Synonyms: HBBM, HBGF-8, NEGF1, HARP

Source: Genetically modified E.coli.

Predicted MW: Monomer, 15.4 kDa (with 137 amino acids)

Presentation: Sterile filtered white lyophilized (freeze-dried) powder. Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5. Lyophilized product is very stable at -20°C. Reconstituted material should be aliquoted and frozen at -20°C to -80°C. It is recommended that a carrier protein (0.1% BSA) is added for long term storage.

Reconstitution: Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. DO NOT VORTEX. Allow several minutes for complete reconstitution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/ml.

Purity determined: Purity is >95% as determined by reducing and non-reducing SDS-PAGE.

Endotoxin Level: Endotoxin level, as measured by LAL analysis, is <0.05 EUs/µg protein (50% confidence range).

AA Sequence: MGKKEKPEKK VKKSDCGEWQ WSVCVPTSGD CGLGTREGTR TGAECKQTMK **TQRCKIPCNW** KKQFGAECKY QFQAWGECDL NTALKTRTGS LKRALHNAEC **OKTVTISKPC** GKLTKPKPQA ESKKKKKEGK KQEKMLD

Laboratory Reagent For Research Use Only

MW 02/15/23

Visit our website for your local distributor.



In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020 e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138 e-mail: service@cedarlanelabs.com

An ISO 9001:2015 and ISO 13485:2016 registered company.