



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Recombinant Human Interferon-gamma (IFN- γ)

CLY100-77-20UG
CLY100-77-100UG
CLY100-77-250UG
CLY100-77-500UG
CLY100-77-1MG

Description: Interferon gamma, IFN γ , is a major immune-modulating molecule produced mainly by T cells and natural killer cells activated by antigens, mitogens or alloantigens. Most immune cells express IFN- γ receptors and respond to IFN γ -induced signaling by up-regulating MHC class I expression. Human IFN γ does not show cross-reactivity with mouse. Recombinant human IFN γ is a non-glycosylated protein, containing 144 amino acids, with a molecular weight of 16.88 kDa.

Source: *E.coli*

Presentation: 20 μ g (**CLY100-77-20UG**), 100 μ g (**CLY100-77-100UG**), 250 μ g (**CLY100-77-250UG**), 500 μ g (**CLY100-77-500UG**) or 1 mg (**CLY100-77-1MG**), sterile filtered white lyophilized (freeze-dried) powder. Recombinant human IFN γ is lyophilized from 5 mM Na₂PO₄, pH 7.2 and 50 mM NaCl.

Reconstitution: Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/mL, which can be further diluted into other aqueous solutions.

Stability: Lyophilized product is very stable at -20°C. Reconstituted material should be aliquoted and frozen at -20°C. It is recommended that a carrier protein (0.1% HSA or BSA) is added for long term storage.

Protein Content and Purity: Typically \geq 95%, determined by HPLC, Reducing and Non-reducing SDS-PAGE, UV spectroscopy at 280 nm.

Biological Activity: The activity is measured in antiviral assays using virally infected L929 cells and is typically less than 0.1 ng/mL.

AA Sequence: MQDPYVKEAE NLKKYFNAGH SDVADNGTLF LGILKNWKEE SDRKIMQSQI
VSFYFKLQFN FKDDQSIQKS VETIKEDMNV KFFNSNKKKR DDFEKLTYNS VTDLNVQRKA
IHELQVMAE LSPAAGTGKR KRSQMLFQGR RASQ

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

References:

1. Interferon-gamma-oligodendrocyte interactions in the regulation of experimental autoimmune encephalomyelitis. *J Neurosci* 2007 Feb 21;27(8):2013-24
2. Role of interferon-gamma in Valpha14+ natural killer T cell-mediated host defense against *Streptococcus pneumoniae* infection in murine lungs. *Microbes Infect* 2007 Jan 9;
3. Blood serum interferon-gamma bioactivity is low in weanling mice subjected to acute deficits of energy or both protein and energy. *Br J Nutr* 2007 Mar;97(3):528-34
4. Pulmonary epithelial cells are a source of interferon-gamma in response to *Mycobacterium tuberculosis* infection. *Immunol Cell Biol* 2007 Feb 20;
5. Effect of Proinflammatory Cytokines, Tumor Necrosis Factor-alpha and Interferon-gamma on Epithelial Barrier Function and Matrix Metalloproteinase-9 in Madin Darby Canine Kidney Cells. *Cell Physiol Biochem* 2007;19(1-4):99-112
6. Inhibition of Intrinsic Interferon-gamma Function Prevents Neointima Formation After Balloon Injury. *Hypertension* 2007 Feb 19.

For Research Use Only

MW 07/19/2019