



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically  
Speaking

**CEDARLANE**<sup>®</sup>  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)



Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

## Recombinant Human Interferon-gamma (IFN- $\gamma$ ) (AF)

**CLY100-77AF-20UG**  
**CLY100-77AF-100UG**  
**CLY100-77AF-250UG**  
**CLY100-77AF-500UG**  
**CLY100-77AF-1MG**

**Description:** Interferon gamma, IFN $\gamma$ , is a major immune-modulating molecule produced mainly by T cells and natural killer cells activated by antigens, mitogens or alloantigens. Most immune cells express IFN- $\gamma$  receptors and respond to IFN $\gamma$ -induced signaling by up-regulating MHC class I expression. Human IFN $\gamma$  does not show cross-reactivity with mouse. Recombinant human IFN $\gamma$  is a non-glycosylated protein, containing 144 amino acids, with a molecular weight of 16.88 kDa.

**Source:** *E.coli*. This product is produced with no animal or human origin raw products. All processing and handling employs animal free equipment and animal free protocols.

**Presentation:** 20  $\mu$ g (CLY100-77AF-20UG), 100  $\mu$ g (CLY100-77AF-100UG), 250  $\mu$ g (CLY100-77AF-250UG), 500  $\mu$ g (CLY100-77AF-500UG) or 1 mg (CLY100-77AF-1MG), sterile filtered white lyophilized (freeze-dried) powder. Recombinant human IFN $\gamma$  is lyophilized from 5 mM Na<sub>2</sub>PO<sub>4</sub>, pH 7.2 and 50 mM NaCl.

**Reconstitution:** Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/mL, which can be further diluted into other aqueous solutions.

**Stability:** Lyophilized product is very stable at -20°C. Reconstituted material should be aliquoted and frozen at -20°C. It is recommended that a carrier protein (0.1% HSA or BSA) is added for long term storage.

**Protein Content and Purity:** Typically  $\geq$  95%, determined by HPLC, Reducing and Non-reducing SDS-PAGE, UV spectroscopy at 280 nm.

**Biological Activity:** The activity is measured in antiviral assays using virally infected L929 cells and is typically less than 0.1 ng/mL.

**AA Sequence:** MQDPYVKEAE NLKKYFNAGH SDVADNGTLF LGILKNWKEE SDRKIMQSQI  
VSFYFKLFDKN FKDDQSIQKS VETIKEDMNV KFFNSNKKKR DDFEKLTYNS VTDLNVQRKA  
IHელიQVMAE LSPAAGTKGR KRSQMLFQGR RASQ

Visit our website for your local distributor.

**CEDARLANE**<sup>®</sup>



[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: [general@cedarlanelabs.com](mailto:general@cedarlanelabs.com)

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: [service@cedarlanelabs.com](mailto:service@cedarlanelabs.com)

**References:**

1. Interferon-gamma-oligodendrocyte interactions in the regulation of experimental autoimmune encephalomyelitis. *J Neurosci* 2007 Feb 21;27(8):2013-24
2. Role of interferon-gamma in Valpha14+ natural killer T cell-mediated host defense against *Streptococcus pneumoniae* infection in murine lungs. *Microbes Infect* 2007 Jan 9;
3. Blood serum interferon-gamma bioactivity is low in weanling mice subjected to acute deficits of energy or both protein and energy. *Br J Nutr* 2007 Mar;97(3):528-34
4. Pulmonary epithelial cells are a source of interferon-gamma in response to *Mycobacterium tuberculosis* infection. *Immunol Cell Biol* 2007 Feb 20;
5. Effect of Proinflammatory Cytokines, Tumor Necrosis Factor-alpha and Interferon-gamma on Epithelial Barrier Function and Matrix Metalloproteinase-9 in Madin Darby Canine Kidney Cells. *Cell Physiol Biochem* 2007;19(1-4):99-112
6. Inhibition of Intrinsic Interferon-gamma Function Prevents Neointima Formation After Balloon Injury. *Hypertension* 2007 Feb 19.

**For Research Use Only**

MW 07/19/2019