



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GFH123

## Recombinant Human GRO-β / CXCL2

### Description

Growth regulated protein β (GRO-β), also known as CXCL2, is a chemokine that is secreted by macrophages, neutrophils, and monocytes at sites of inflammation. GRO-β functions as a chemoattractant for leukocytes and hematopoietic stem cells. GRO-β activity is mediated through binding the G-protein-coupled chemokine receptor CXCR2.

<b>Length</b>	73 aa
<b>Molecular Weight</b>	7.9 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	P19875
<b>Purity</b>	≥95% determined by reducing and non-reducing SDS-PAGE


### Specifications

<b>Alternative Names</b>	Growth regulated protein β, GROb, GROb1, CXCL2, MGSAβ, MGSA β, MGSA-b, CXCL2, MIP-2 α, MIP-2a, MIP2-α, MIP2A, MIP22, GRO2, chemokine (C-X-C motif) ligand 2, C-X-C motif chemokine 2, SCYB2, melanoma growth stimulatory activity β, CINC-2a, GROB3, GRO2 oncogene, macrophage inflammatory protein 2 α
<b>Biological Activity</b>	Human GRO-β is fully biologically active when compared to standard. The activity is determined by the induced chemotaxis of human neutrophils.
<b>Endotoxin Level</b>	≤1.00 EU/μg as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
<b>AA Sequence</b>	APLATELRQC CLQTLQGIHL KNIQSVKVKVKS PGPHCAQTEV IATLKNGQKA CLNPASPMVK KIIEKMLKNG KSN


### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
<b>Stability and Storage</b>	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

### Data



Induced chemotaxis of human neutrophils assay for Human GRO-β.



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μg of protein was loaded in each lane. Human GRO-β has a predicted Mw of 7.9 kDa.