



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GFH128 Recombinant Human ITAC / CXCL11

### Description

Interferon-inducible T cell  $\alpha$  chemoattractant (ITAC), also known as CXCL11, is expressed at high levels in leukocytes, pancreas, and liver cells. ITAC gene expression is induced by interferons  $\alpha$  (IFN- $\alpha$ ),  $\beta$  (IFN- $\beta$ ), and  $\gamma$  (IFN- $\gamma$ ). ITAC is the dominant ligand known to bind the chemokine receptor CXCR3, thus acting as a strong agonist. ITAC functions as a chemoattractant for interleukin-2 (IL-2)-activated T cells.

<b>Length</b>	73 aa
<b>Molecular Weight</b>	8.3 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	O14625
<b>Purity</b>	$\geq$ 95% determined by reducing and non-reducing SDS-PAGE


### Specifications

<b>Alternative Names</b>	Interferon-inducible T cell $\alpha$ chemoattractant, CXCL11, B-R1, I-TAC, chemokine (C-X-C motif) ligand 11, C-X-C motif chemokine 11
<b>Biological Activity</b>	Human ITAC is fully biologically active when compared to standard. The activity is determined by the ability to chemoattract IL-2 activated T cells at a concentration of 0.1 - 10 ng/ml. There is no data currently available.
<b>Endotoxin Level</b>	$\leq$ 1.00 EU/ $\mu$ g as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
<b>AA Sequence</b>	FPMFKRGRCL CIGPGVKAVK VADIEKASIM YPSNNDKIE VIITLKENKG QRCLNPKSKQ ARLIKKVER KNF

### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
<b>Stability and Storage</b>	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

### Data



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1  $\mu$ g of protein was loaded in each lane. Human ITAC has a predicted Mw of 8.3 kDa.