



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GFH147 Recombinant Human MIP-3 $\alpha$ / CCL20

### Description

Macrophage Inflammatory Protein 3  $\alpha$  (MIP-3  $\alpha$ ), also known as CCL20, is expressed in the liver, lungs, lymph nodes, and peripheral blood lymphocytes. MIP-3  $\alpha$  expression is strongly induced by inflammatory signals, and downregulated by the anti-inflammatory cytokine interleukin-10 (IL-10). MIP-3  $\alpha$  signals through the G protein-coupled receptor CCR6 to function as a chemoattractant to lymphocytes and dendritic cells.

<b>Length</b>	70 aa
<b>Molecular Weight</b>	8.0 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	P78556
<b>Purity</b>	≥95% determined by reducing and non-reducing SDS-PAGE


### Specifications

<b>Alternative Names</b>	Monocyte Chemotactic Protein 3, MIP-3 alpha, CCL20
<b>Biological Activity</b>	Human MIP-3 $\alpha$ is fully biologically active when compared to standard. The activity is determined by the ability to induce chemotaxis of primary human T cells.
<b>Endotoxin Level</b>	≤1.00 EU/ $\mu$ g as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
<b>AA Sequence</b>	ASNFDCCCLGY TDRILHPKFI VGFTRQLANE GCDINAIIFH TKKKLSVCAN PKQTWVKYIV RLLSKKVKNM

### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
<b>Stability and Storage</b>	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

### Data



Induced chemotaxis of human primary T cells assay for Human MIP-3  $\alpha$ . Cells that migrated were counted using a luminescent substrate. Migration over basal levels was reported in response to Human MIP-3  $\alpha$  starting at 5 ng/ml.