



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFH170

Recombinant Human GRO- γ / CXCL3

Description

Growth regulated protein γ (GRO- γ), also called CXCL3, acts through the chemokine receptor CXCR2 to promote monocyte migration and adhesion. GRO- γ also induces the migration of cerebellar granule neuron precursor cells.

Length	73 aa
Molecular Weight	7.9 kDa
Source	E. coli
Accession Number	P19876
Purity	\geq 95% determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Growth regulated protein β , GRO β , GRO β 1, CXCL2, MGSA β , MGSA β , MGSA-b, CXCL2, MIP-2 α , MIP-2a, MIP2- α , MIP2A, MIP22, GRO2, chemokine (C-X-C motif) ligand 2, C-X-C motif chemokine 2, SCYB2, melanoma growth stimulatory activity β , CINC-2a, GROB3, GRO2 oncogene, macrophage inflammatory protein 2 α
Biological Activity	Human GRO- γ is fully biologically active when compared to standard. The activity is determined by the induced chemotaxis of human neutrophils.
Endotoxin Level	\leq 1.00 EU/ μ g as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	ASVVTELRCQ CIQLTQGIHL KNIQSVNVRS PGPHCAQTEV IATLKNKGKA CLNPASPMVQ KIEKILNKG STN


Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Induced chemotaxis of human neutrophils assay for Human GRO- γ .



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μ g of protein was loaded in each lane. Human GRO- γ has a predicted Mw of 7.9 kDa.