



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GFH182AF Recombinant Human IHH (Animal-Free)

### Description

Indian hedgehog (IHH) is an essential signaling factor that is secreted in the gut, cartilage, and bone during embryonic development. IHH acts through the patched (PTC) receptor to induce transcriptional changes important for bone and cartilage development. IHH also induces the expression of parathyroid hormone-related peptide (PTHrP), which in turn mediates IHH activity during chondrocyte differentiation, forming a negative feedback loop.

This product is produced with no animal derived raw products. All processing and handling employs animal free equipment and animal free protocols.

<b>Length</b>	177 aa
<b>Molecular Weight</b>	19.9 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	Q14623
<b>Purity</b>	≥95% determined by reducing and non-reducing SDS-PAGE


### Specifications

<b>Alternative Names</b>	Indian Hedgehog, HHG-2
<b>Biological Activity</b>	Human IHH (Animal-Free) is fully biologically active when compared to standard. The activity is determined by the induced alkaline phosphatase activity of CCL-26 cells.
<b>Endotoxin Level</b>	≤1.00 EU/μg as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
<b>AA Sequence</b>	MIIGPGRVVG SRRRPPrKLV PLAYKQFSPN VPEKTLGASG RYEGKIARSS ERFKELTPNY NPDIIIFKDEE NTGADRLMTQ RCKDRLNSLA ISVMNQWPGV KLRVTEGWDE DGHHSSESLH YEGRAVDITT SDRDRNKYGL LARLAVEAGF DWVYYESKAH VHCSVKSEHS AAAKTGG

### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile 10 mM HCl at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
<b>Stability and Storage</b>	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

### Data



Induced alkaline phosphatase activity of CCL-26 cells assay for Human IHH. Cell proliferation was measured to calculate the ED50, which is in the range of μg/ml.