



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFH185 Recombinant Human ENA-78 / CXCL5

Description

Epithelial-derived neutrophil-activating peptide 78 (ENA-78), also known as CXCL5, is a chemokine that recruits neutrophils, promotes angiogenesis, and stimulates connective tissue remodelling. ENA-78 production is stimulated by Interleukin-1 (IL-1) or tumor Necrosis Factor α (TNF- α), and signals through the chemokine receptor CXCR2. ENA-78 with 5 - 78 aminoacids is one of three naturally occurring ENA-78 variants in which the N-terminus has been truncated.

Length	74 aa
Molecular Weight	8.1 kDa
Source	E. coli
Accession Number	P42830
Purity	\geq 95% determined by reducing and non-reducing SDS-PAGE

Specifications

Alternative Names	CXCL5, epithelial-derived neutrophil-activating protein 78, ENA78, ENA 78
Biological Activity	Human ENA-78 is fully biologically active when compared to standard. The activity is determined by the ability to induce chemotaxis of primary human neutrophils starting at a concentration of 10 ng/ml. This corresponds to an expected specific activity of 1×10^5 units/mg. There is no data currently available.
Endotoxin Level	\leq 1.00 EU/ μ g as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	AAVLRELRCV CLQTTQGVHP KMISNLQVFA IGPQCSKVEV VASLKNGKEI CLDPEAPFLK KVIQKILDGG NKEN

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.