



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFH32

Recombinant Human GRO- α / CXCL1

Description

Growth regulated protein α (GRO- α), also known as CXCL1, is a chemokine that has mitogenic properties and is a neutrophil chemoattractant. GRO- α is secreted by macrophages, epithelial cells, neutrophils, and melanomas. GRO- α signals through the CXCR2 chemokine receptor and is important during spinal cord formation, inflammation, angiogenesis, tumorigenesis, and wound healing.

| | |
|-------------------------|--|
| Length | 73 aa |
| Molecular Weight | 7.9 kDa |
| Source | E. coli |
| Accession Number | P09341 |
| Purity | $\geq 95\%$ determined by reducing and non-reducing SDS-PAGE |


Specifications

| | |
|----------------------------|--|
| Alternative Names | Growth regulated protein α , GRO α , GRO α |
| Biological Activity | Human GRO- α is fully biologically active when compared to standard. The activity is determined by the induced chemotaxis of human neutrophils. |
| Endotoxin Level | ≤ 1.00 EU/ μ g as measured by kinetic LAL |
| Formulation | Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA) |
| AA Sequence | ASVATELRQC CLQTLQGIHP KNIQSVNVKS PGPCHAQTEV IATLKNGRKA CLNPASPIVK KIIIEKMLNSD KSN |


Preparation and Storage

| | |
|------------------------------|---|
| Reconstitution | Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions. |
| Stability and Storage | 12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed. |

Data



Induced chemotaxis of human neutrophils assay for Human GRO- α .



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μ g of protein was loaded in each lane. Human GRO- α has a predicted Mw of 7.9 kDa.