



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFH52 Recombinant Human IFN- α 2b

Description

Interferon- α 2b (IFN- α 2b) is a type I interferon made by leukocytes during viral infection. The JAK-STAT pathway mediates the antiviral and anti-cell proliferation activities of IFN- α 2b. IFN- α proteins are widely used as standard treatments during antiviral and antineoplastic therapies. The IFN- α 2b variant differs from IFN- α 2b by one amino acid.

Length	166 aa
Molecular Weight	19.4 kDa
Source	E. coli
Accession Number	P01563 (VAR_004012)
Purity	\geq 95% determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Interferon- α 2b, interferon alpha, interferon-alpha, type I Interferon α 2, IFN α 2
Biological Activity	Human IFN- α 2b is fully biologically active when compared to standard. The activity is determined by the viral CPE assay using EMC virus on A549 cells. The activity corresponds to an expected specific activity of 2.0×10^8 units/mg.
Endotoxin Level	\leq 1.00 EU/ μ g as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	MCDLPQTHSL GSRRTLMLLA QMRRISLFSC LKDRHDFGFP QEEFGNQFQK AETIPVLHEM IQQIFNLFST KDSSAAWDET LLDKPHYTELY QQLNDLEACV IQGVGVTEPTP LMKEDSILAV RKYFQRITLY LKEKKYSPCA WEVVRAEIMR SFSLSLSTNLQE SLRSKE


Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Viral CPE assay using EMC virus on A549 cells for Human IFN- α 2b.



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μ g of protein was loaded in each lane. Human IFN- α 2b has a predicted Mw of 19.4 kDa.