



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GFM12 Recombinant Mouse FGF-2

### Description

Fibroblast Growth Factor 2 (FGF-2) is expressed by endothelial cells and is a mediator of angiogenesis. FGF-2 also has cardioprotective functions during heart injury. The application of FGF-2 is a critical component for human embryonic stem cell culture systems and is necessary for maintaining human embryonic stem cells in an undifferentiated state.

<b>Length</b>	146 aa
<b>Molecular Weight</b>	16.5 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	P15655
<b>Purity</b>	≥95% determined by reducing and non-reducing SDS-PAGE

### Specifications

<b>Alternative Names</b>	Fibroblast Growth Factor 2, FGF2, FGF 2, HBGF-2, basic fibroblast growth factor, heparin-binding growth factor 2, FGFB, BFGF, bFGF, prostatropin
<b>Biological Activity</b>	Mouse FGF-2 is fully biologically active when compared to standard. The activity is determined by the dose-dependent induced proliferation of NR6R 3T3 cells and it is typically less than 2.5 ng/ml. This corresponds to an expected specific activity of 4.0 x 10 <sup>5</sup> units/mg.
<b>Endotoxin Level</b>	≤1.00 EU/μg as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5
<b>AA Sequence</b>	MPALPEDGGA AFPPGHFKDP KRLYCKNGGF FLRIHPDGRV DGVREKSDPH VKLQLQAEER GVVSIKGVCA NRYLAMKEDG RLLASKCVTE ECVFERLES NNYNTYRSRK YSSWYVALKR TGQYKLGSKT GPGQKAILFL PMSAKS

### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.
<b>Stability and Storage</b>	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.