



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFM12AF Recombinant Mouse FGF-2 (Animal-Free)

Description

Fibroblast Growth Factor 2 (FGF-2) is expressed by endothelial cells and is a mediator of angiogenesis. FGF-2 also has cardioprotective functions during heart injury. The application of FGF-2 is a critical component for human embryonic stem cell culture systems and is necessary for maintaining human embryonic stem cells in an undifferentiated state.

This product is produced with no animal derived raw products. All processing and handling employs animal free equipment and animal free protocols.

Length	146 aa
Molecular Weight	16.5 kDa
Source	E. coli
Accession Number	P15655
Purity	≥95% determined by reducing and non-reducing SDS-PAGE

Specifications

Alternative Names	Fibroblast Growth Factor 2, FGF2, FGF 2, HBGF-2, basic fibroblast growth factor, heparin-binding growth factor 2, FGFB, BFGF, bFGF, prostatropin
Biological Activity	Mouse FGF-2 is fully biologically active when compared to standard. The activity is determined by the dose-dependent induced proliferation of NR6R 3T3 cells and it is typically less than 2.5 ng/ml. This corresponds to an expected specific activity of 4.0×10^5 units/mg.
Endotoxin Level	≤1.00 EU/μg as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5
AA Sequence	MPALPEDGGA AFPPGHFKDP KRLYCKNGGF FLRIHPDGRV DGVREKSDPH VKLQLQAEER GVVSIKGVCA NRYLAMKEDG RLLASKCVTE ECFFERLES NNYNTYRSRK YSSWYVALKR TGQYKLGSKT GPGQKAILFL PMSAKS

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.