



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFM16AF Recombinant Mouse IFN- γ (Animal-Free)

Description

Interferon γ (IFN- γ) is a type II interferon that is critical during adaptive and innate immune responses to infection. IFN- γ is produced by T cells and Natural Killer cells following antigen-specific activation. IFN- γ binds IFN- γ receptors (IFN- γ R1 and IFN- γ R2), which are expressed on most immune cells, to activate the JAK-STAT pathway. IFN- γ induced signaling increases the expression of class I major histocompatibility complex (MHC) molecules. Mouse IFN- γ is not cross-reactive with human IFN- γ .

This product is produced with no animal derived raw products. All processing and handling employs animal free equipment and animal free protocols.

Length	134 aa
Molecular Weight	15.7 kDa
Source	E. coli
Accession Number	P01580
Purity	$\geq 95\%$ determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Interferon- γ , interferon-gamma, interferon gamma, type II interferon, T cell interferon, immune Interferon, MAF, IFG, IFI
Biological Activity	Mouse IFN- γ is fully biologically active when compared to standard. The activity is determined by the viral CPE assay using EMC virus on L929 cells. The activity corresponds to an expected specific activity of 1.0×10^7 units/mg.
Endotoxin Level	≤ 1.00 EU/ μ g as measured by kinetic LAL
Formulation	Lyophilized from a sterile filtered solution containing 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5
AA Sequence	MHGTVIESLE SLNNYFNSSG IDVEEKSLFL DIWRNWQKDG DMKILQSQII SFYLRLFEVL KDNQAISNNI SVIESHLITT FFSNSKAKKD AFMSIAKFEV NNPQVQRQAF NELIRVVHQL LPESLRLKRRK RSRC


Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions. If a precipitate is observed, centrifuge the solution thoroughly and use only the soluble fraction (removing it from the precipitate). A 10% overfill has been added to compensate for any loss of protein in the precipitate.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Viral CPE assay using EMC virus on L929 cells for Mouse IFN- γ .



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μ g of protein was loaded in each lane. Mouse IFN- γ has a predicted Mw of 15.7 kDa.