



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GFM24 Recombinant Mouse Follistatin

### Description

Follistatin is an autocrine, activin-binding protein that is ubiquitously expressed with highest expression levels being in the ovary and skin. Follistatin negatively regulates the signaling of Transforming Growth Factor  $\beta$  (TGF- $\beta$ ) family members, such as activin, bone morphogenic proteins (BMP), myostatin, Growth Differentiation Factor 11 (GDF-11), and TGF- $\beta$ 1. Follistatin functions as an antagonist by binding TGF- $\beta$  family members to block interaction with their signaling receptors. Follistatin also inhibits the secretion of Follicle-Stimulating Hormone (FSH) from the anterior pituitary.

<b>Length</b>	289 aa
<b>Molecular Weight</b>	31.7 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	P19883
<b>Purity</b>	$\geq$ 95% determined by reducing and non-reducing SDS-PAGE


### Specifications

<b>Alternative Names</b>	FS, activin-binding protein, FSH-suppressing protein, FSP
<b>Biological Activity</b>	Mouse Follistatin is fully biologically active when compared to standard. The activity is determined by the neutralization of Activin A.
<b>Endotoxin Level</b>	$\leq$ 1.00 EU/ $\mu$ g as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5
<b>AA Sequence</b>	<p>MGNCWLRQAK NGRCQVLYKT ELSKEECCST GRLSTSWTEE DVNDNTLFWK MIFNGGAPNC</p> <p>IPCKETCENV DCGPGKRCRM NKKNKPRVC APDCSNITWK GPVCGLDGKT YRNECALLKA</p> <p>RCKEQPELEV QYQGRCKKTC RDVFCPGSST CVVDQTNNAY CVTCNRICPE PASSEQYLCG</p> <p>NDGVTYSSAC HLRKATCLLG RSLGLAYEGK CIKAKSCEDI QCTGGKKCLW DF</p>

### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
<b>Stability and Storage</b>	<p>12 months from date of receipt when stored at -20°C to -80°C as supplied.</p> <p>1 month when stored at 4°C after reconstituting as directed.</p> <p>3 months when stored at -20°C to -80°C after reconstituting as directed.</p>

### Data



Neutralization of Activin A induced cytotoxicity of MPC-11 cells assay for Mouse Follistatin. Cell proliferation was measured to calculate the ED50.