



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GFM38

## Recombinant Mouse IP-10 / CXCL10

### Description

Interferon  $\gamma$ -induced protein 10 (IP-10), also known as CXCL10, is a chemokine secreted by monocytes, endothelial cells and fibroblasts in response to interferon  $\gamma$  (IFN- $\gamma$ ). IP-10 functions as a chemoattractant for activated T cells, monocytes, dendritic, and Natural Killer (NK) cells that express the G protein-coupled receptor CXCR3. IP-10 is an important factor in autoimmune diseases such as Hashimoto's thyroiditis, Graves' disease, and Type 1 diabetes mellitus.

<b>Length</b>	76 aa
<b>Molecular Weight</b>	8.7 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	P13500
<b>Purity</b>	$\geq$ 95% determined by reducing and non-reducing SDS-PAGE


### Specifications

<b>Alternative Names</b>	Monocyte Chemotactic Protein 1, CCL2, JE, MCAF
<b>Biological Activity</b>	Mouse IP-10 is fully biologically active when compared to standard. The activity is determined by the ability to induce chemotaxis of primary human T cells.
<b>Endotoxin Level</b>	$\leq$ 1.00 EU/ $\mu$ g as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
<b>AA Sequence</b>	QPDAINAPVT CCYNFTNRKI SVQRLASYRR ITSSKCPKEA VIFKTIVAKE ICADPKQKWV QDSMDHLDKQ TQTPKT

### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
<b>Stability and Storage</b>	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

### Data



Induced chemotaxis of primary human T cells assay for Mouse IP-10. Cells that migrated were counted using a luminescent substrate. Migration over basal levels was reported in response to Mouse IP-10 starting at 5 ng/ml.