



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFM3AF Recombinant Mouse IL-7 (Animal-Free)

Description

Interleukin-7 (IL-7) is a hematopoietic cytokine that is an important regulator of B and T cell development. IL-7 is secreted by bone marrow and thymic stromal cells, dendritic cells, intestinal epithelial cells, hepatocytes, and keratinocytes. IL-7 signals through the interleukin-7 receptor (IL-7R) to promote the differentiation of hematopoietic stem cells into T cells, B cells, and Natural Killer cells. IL-7 is also a regulator of intestinal mucosal lymphocyte proliferation. Human and mouse IL-7 show species cross-reactivity.

This product is produced with no animal derived raw products. All processing and handling employs animal free equipment and animal free protocols.

Length	130 aa
Molecular Weight	15 kDa
Source	E. coli
Accession Number	P10168
Purity	≥95% determined by reducing and non-reducing SDS-PAGE

Specifications

Alternative Names	Interleukin-7, interleukin 7, IL7, IL 7, lymphopoietin 1, LP-1, pre-B cell factor
Biological Activity	Mouse IL-7 is fully biologically active when compared to standard. The activity is determined by the ability to induce mouse 2E8 cells proliferation and it is typically less than 1 ng/ml. This corresponds to an expected specific activity of 1×10^6 units/mg.
Endotoxin Level	≤1.00 EU/μg as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	MECHIKDKEG KAYESVLMIS IDELDKMTGT DSNCPNNEPN FFRKHVCDT KEAAFLNRAA RCLKQFLKMN ISEEFNVHLL TVSQGTQTLV NCTSKEEKNV KEQKNDACF LKRLLEIKT CWNKILKGS I

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile 10 mM acetic acid at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.