

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





DATA SHEET

GFM40

Recombinant Mouse MCP-3 / CCL7

Description

Monocyte Chemotactic Protein 3 (MCP-3), also known as CCL7, is produced by macrophages and tumor cell lines. MCP-3 signals through the G protein-coupled receptors CCR1, CCR2, and CCR3. MCP-3 chemoattracts monocytes and regulates macrophage function during inflammation and metastasis.

Length74 aaMolecular Weight8.5 kDaSourceE. coliAccession NumberQ03366

Purity ≥95% determined by reducing and non-reducing SDS-PAGE

Specifications

Alternative Names Monocyte Chemotactic Protein 1, CCL2, JE, MCAF

Biological Activity Mouse MCP-3 is fully biologically active when compared to standard. The activity is determined by the ability

to induce chemotaxis of THP-1 cells.

Endotoxin Level ≤1.00 EU/µg as measured by kinetic LAL

Formulation Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)

AA Sequence QPDGPNASTC CYVKKQKIPK RNLKSYRRIT SSRCPWEAVI FKTKKGMEVC AEAHQKWVEE

AIAYLDMKTP TPKP

Preparation and Storage

Reconstitution Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the

vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized


product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.

Stability and Storage 12 months from date of receipt when stored at -20°C to -80°C as supplied.

1 month when stored at 4°C after reconstituting as directed.

3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Induced chemotaxis of THP-1 cells assay for Mouse MCP-3. Cells that migrated were counted using a luminescent substrate. Migration over basal levels was reported in response to Mouse MCP-3 starting at 1000 ng/ml.