



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFM52 Recombinant Mouse VEGF-120

Description

Vascular Endothelial Growth Factor 120 (VEGF-120) is produced by a wide variety of cell types, including tumor and vascular cells. VEGF-120 is a mediator of vascular growth, vascular permeability, and plays a role in stimulating vasodilation via nitric oxide-dependent pathways. Mouse VEGF-A has several alternatively spliced isoforms, with one being VEGF-120. VEGF-120 is an angiogenic factor that is expressed throughout endochondral bone development and is important during skeletogenesis.

Length	121 / 242 aa
Molecular Weight	14.1 / 28.3 kDa
Source	E. coli
Accession Number	P15692-9
Purity	≥95% determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Vascular Endothelial Growth Factor, VEGF121, VEGF
Biological Activity	Mouse VEGF-120 is fully biologically active when compared to standard. The activity is determined by the proliferation of HUVEC cells and it is typically less than 5 ng/ml. This corresponds to an expected specific activity of 2.0×10^5 units/mg.
Endotoxin Level	≤1.00 EU/μg as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	MPMAEGGGQN HHEVVKFMDV YQRSYCHPIE TLVDIFQEYP DEIEYIFKPS CVPLMRCGGC CNDEGLECVP TEESNITMQI MRIKPHQGQH IGEMSFLLQHN KCECRPKKDR ARQENCDKPR R

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Induced proliferation of HUVEC cells for Mouse VEGF-120. Cell proliferation was measured to calculate the ED50.