



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GFM54 Recombinant Mouse PDGF-AA

### Description

Platelet-Derived Growth Factor (PDGF) is an important regulator of cell growth, proliferation, and angiogenesis. PDGF synthesis is induced by IL-1, IL-6, TNF- $\alpha$ , TGF- $\beta$  and EGF signaling. PDGF functions as a mitogenic growth hormone on cells of mesenchymal lineage, such as smooth muscle and glial cells. PDGF is also stored in the  $\alpha$ -granules of platelets and is released upon adherence to traumatized tissues. PDGF is a dimeric glycoprotein formed by two A chains (AA), two B chains (BB), or as a heterodimer with an A and a B chain (AB). The PDGF dimer binds the cell surface receptor tyrosine kinases PDGFR-a and PDGFR-b.

<b>Length</b>	126 / 252 aa
<b>Molecular Weight</b>	14.4 / 28.9 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	P04085
<b>Purity</b>	$\geq 95\%$ determined by reducing and non-reducing SDS-PAGE


### Specifications

<b>Alternative Names</b>	Platelet-Derived Growth Factor, GDGF, ODGF, PDGF AA
<b>Biological Activity</b>	Mouse PDGF-AA is fully biologically active when compared to standard. The activity is determined by the induction of BALB/c 3T3 cells proliferation and it is typically less than 10 ng/ml. This corresponds to an expected specific activity of $1.0 \times 10^5$ units/mg.
<b>Endotoxin Level</b>	$\leq 1.00$ EU/ $\mu$ g as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
<b>AA Sequence</b>	MSIEEAVPAV CKTRTVIYEI PRSQVDPTSA NFLIWPPCVE VKRCTGCCNT SSVKQPSRV HHRSVKVAKV EYVRKKPKLK EVQVRLEEHL ECACATTSLN PDYREEDTGR PRESGKKRKR KRLKPT

### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
<b>Stability and Storage</b>	12 months from date of receipt when stored at $-20^{\circ}\text{C}$ to $-80^{\circ}\text{C}$ as supplied. 1 month when stored at $4^{\circ}\text{C}$ after reconstituting as directed. 3 months when stored at $-20^{\circ}\text{C}$ to $-80^{\circ}\text{C}$ after reconstituting as directed.

### Data



Induced proliferation of BALB/c 3T3 cells assay for Mouse PDGF-AA. Cell proliferation was measured to calculate the ED50, which is as expected less than 10 ng/ml.