



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



GFM62

Recombinant Mouse IL-17F

Description

Interleukin-17F (IL-17F), a member of the IL-17 cytokine family, is secreted by activated CD4+ T cells and monocytes. IL-17F binds the IL-17 receptor related molecule, IL17RC, to promote the production of the interleukin-6 (IL-6), interleukin-8 (IL-8), and Granulocyte Macrophage Colony-Stimulating Factor (GM-CSF) cytokines. IL-17F also functions to regulate matrix turnover rates, inhibit endothelial cell angiogenesis, and induce the endothelial cell expression of interleukin-2 (IL-2), Monocyte Chemoattractant Protein 1 (MCP-1), and Transforming Growth Factor β 1 (TGF- β 1).

Length	134 / 268 aa
Molecular Weight	15.0 / 30.0 kDa
Source	E. coli
Accession Number	Q7TNI7
Purity	$\geq 95\%$ determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Interleukin-17F, interleukin 17F, IL17F, IL 17F
Biological Activity	Mouse IL-17F is fully biologically active when compared to standard. The activity is determined by the ability to produce mouse IL-6 by NIH-3T3 cells.
Endotoxin Level	≤ 1.00 EU/ μ g as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	MRKNPKAGVP ALQKAGNCPP LEDNTVRVDI RIFNQNQGIS VPREFQRSS SPWDYNITRD PHRFPSAE AQCRHSGCIN AQGQEDSTMN SVAIQQEILV LRREPQGCSN SFRLAKMLLK VGCTCVKPIV HQAA

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Induced production of mouse IL-6 by NIH-3T3 cells. Production of mouse IL-6 was measured to calculate the ED50.