



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFM67

Recombinant Mouse SDF-1 β / CXCL12b

Description

Stromal cell-derived factor-1 β (SDF-1 β), also known as CXCL12b, is one of two SDF-1 splice variants made by a wide variety of cells upon stimulation by inflammatory cytokines such as TNF, IL-1, and LPS. SDF-1 β signals through the G protein-coupled receptor CXCR4 to recruit activated leukocytes.

Length	68 aa
Molecular Weight	8.0 kDa
Source	E. coli
Accession Number	P48061 (2)
Purity	\geq 95% determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Monocyte Chemotactic Protein 1, CCL2, JE, MCAF
Biological Activity	Mouse SDF-1 β is fully biologically active when compared to standard. The activity is determined by the ability to induce chemotaxis of human neutrophils.
Endotoxin Level	\leq 1.00 EU/ μ g as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	KPVSLSYRCP CRFFESHVAR ANVKHLKILN TPNCALQIVA RLKNNNRQVC IDPKLKWIQE YLEKALNK

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Induced chemotaxis of neutrophils assay for Mouse SDF-1 β . Cells that migrated were counted using a luminescent substrate. Migration over basal levels was reported in response to Mouse SDF-1 β starting at 1000 ng/ml.