



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFM69

Recombinant Mouse RELM- β

Description

Resistin-like molecule- β (RELM- β) is a member of the RELM-family of secreted proteins containing conserved C-terminus cysteines. The RELM-family consists of resistin (FIZZ3), RELM- α (FIZZ1), RELM- β (FIZZ2), and RELM- γ (FIZZ4). Resistin and RELM- β are the only RELM-family members found in humans, whereas all four RELM-family members are present in rodents. RELM- β functions to increase fibroblast proliferation and differentiation, resulting in airway remodelling and increased inflammation.

Length	83 aa
Molecular Weight	8.9 kDa
Source	E. coli
Accession Number	Q99P86
Purity	\geq 95% determined by reducing and non-reducing SDS-PAGE

Specifications

Alternative Names	Resistin-like molecule- β , RELM β , FIZZ2
Biological Activity	There is no data currently available.
Endotoxin Level	\leq 1.00 EU/ μ g as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	MQCSFESLVD QRIKEALSRQ EPKTISCTSV TSSGRLASCP AGMVVTGCAC GYGCGSWDIR NGNTCHCQCS VMDWASARCC RMA

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.