



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFM70AF Recombinant Mouse IL-21 (Animal-Free)

Description

Interleukin-21 (IL-21) is a member of the common- γ chain family of cytokines that are involved in immunoregulation. IL-21 is normally expressed by activated CD4+ T cells and is aberrantly expressed in Hodgkin lymphoma cells. The IL-21 receptor (IL-21R) activates the JAK/STAT signaling pathway and is expressed on T, B, and Natural Killer (NK) cells. Within the B cell lineage, IL-21 is a switch factor regulating IgG1 and IgG3 antibody production. IL-21 also cooperates with IL-4 for the production of multiple antibody classes in B cells. IL-21 has pleiotropic effects on the proliferation, differentiation, and effector functions of B, T, NK, and dendritic cells.

This product is produced with no animal derived raw products. All processing and handling employs animal free equipment and animal free protocols.

| | |
|-------------------------|---|
| Length | 132 aa |
| Molecular Weight | 15.4 kDa |
| Source | E. coli |
| Accession Number | Q9HBE4 |
| Purity | ≥95% determined by reducing and non-reducing SDS-PAGE |

Specifications

| | |
|----------------------------|---|
| Alternative Names | Interleukin-21, interleukin 21, IL21, IL 21 |
| Biological Activity | Mouse IL-21 is fully biologically active when compared to standard. The activity is determined by the ability to induce B9 cells proliferation and it is typically less than 20 ng/ml. This corresponds to an expected specific activity of 5×10^4 units/mg. |
| Endotoxin Level | ≤1.00 EU/μg as measured by kinetic LAL |
| Formulation | Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA) |
| AA Sequence | MQDRHMIRMRLQLIDIVDQLK NYVNDLVPEFLPAPEDVETN CEWSAFSCFQ KAQLKSANTG NNERIINVSI KKLKRKPPST NAGRRQKHRL TCPSCDSYEK KPPKEFLERF KSLQKMIHQ HLSSRTHGSE DS |

Preparation and Storage

| | |
|------------------------------|---|
| Reconstitution | Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile 10 mM HCl at 0.1 mg/ml, which can be further diluted into other aqueous solutions. |
| Stability and Storage | 12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed. |