



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFM77AF Recombinant Mouse IL-22 (Animal-Free)

Description

Interleukin-22 (IL-22) is an IL-10 family member that is produced by activated dendritic cells and T lymphocytes. IL-22 signals via a heteroduplex receptor consisting of IL-22R and IL-10R β chains. IL-22 is a potent mediator of cellular inflammatory responses.

This product is produced with no animal derived raw products. All processing and handling employs animal free equipment and animal free protocols.

Length	147 aa
Molecular Weight	16.8 kDa
Source	E. coli
Accession Number	Q9JJY9
Purity	≥95% determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Interleukin-22, interleukin 22, IL22, IL 22, IL-TIF
Biological Activity	Mouse IL-22 (Animal-Free) is fully biologically active when compared to standard. The activity is determined by the ability to induce production of IL-10 from human COLO-205 cells and it is typically less than 1 ng/ml. This corresponds to an expected specific activity of 1×10^6 units/mg.
Endotoxin Level	≤1.00 EU/ μ g as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	MLPVTNRCKL EVSNFQQPYI VNRTFMLAKE ASLADNNTDV RLIGEKLFGRG VSAKDQCYLM KQVLNFTLED VLLPQSDRFQ PYMQEVVFFL TKLSNQLSSC HISGDDQNIQ KNVRLKQTV KKLGESGEIK AIGELDLLFM SLRNACV


Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Induced production of IL-10 by human COLO-205 cells. Production of IL-10 was measured to calculate the ED50, which is as expected less than 1 ng/ml.



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μ g of protein was loaded in each lane. Mouse IL-22 has a predicted Mw of 16.8 kDa.