



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GFM78AF Recombinant Mouse MIF (Animal-Free)

### Description

Migration Inhibitory Factor (MIF) is a pro-inflammatory lymphokine that functions during cell-mediated immunity. MIF promotes fibroblast migration by inducing interleukin-1 (IL-1), interleukin-8 (IL-8), and matrix metalloproteinase (MMP) expression. In interferon- $\gamma$ -activated macrophages, MIF stimulates nitric oxide (NO) production and Tumor Necrosis Factor  $\alpha$  (TNF- $\alpha$ ) secretion.

This product is produced with no animal derived raw products. All processing and handling employs animal free equipment and animal free protocols.

<b>Length</b>	115 aa
<b>Molecular Weight</b>	12.5 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	P34884
<b>Purity</b>	$\geq 95\%$ determined by reducing and non-reducing SDS-PAGE

### Specifications

<b>Alternative Names</b>	Migration Inhibitory Factor, GIF, phenylpyruvate tautomerase, glycosylation-inhibiting factor, L-dopachrome tautomerase
<b>Biological Activity</b>	Activity to be determined.
<b>Endotoxin Level</b>	$\leq 1.00$ EU/ $\mu$ g as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5
<b>AA Sequence</b>	MPMFIVNTNV PRASVPEGFL SELTQQLAQA TGKPAQYIAV HVVPDQLMTF SGTNDPCALC SLHSIGKIGG AQNRNYSKLL CGLLSDRLHI SPDRVYINYY DMNAANVGWN GSTFA

### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
<b>Stability and Storage</b>	12 months from date of receipt when stored at $-20^{\circ}\text{C}$ to $-80^{\circ}\text{C}$ as supplied. 1 month when stored at $4^{\circ}\text{C}$ after reconstituting as directed. 3 months when stored at $-20^{\circ}\text{C}$ to $-80^{\circ}\text{C}$ after reconstituting as directed.